

Neutrino Oscillations: a source of Goldstone fields and consequences for Supernovae*

Luís Bento

*Centro de Física Nuclear da Universidade de Lisboa,
Av. Prof. Gama Pinto 2, 1699 Lisboa - codex, Portugal
e-mail: lbento@fc.ul.pt*

Abstract

It is shown that true Nambu-Goldstone (NG) bosons develop coherent fields whenever the associated charges of the matter particles are not conserved in a macroscopic scale. The sources of the NG fields are the time rates of quantum number violation. If the lepton numbers are spontaneously broken at a scale below 1 TeV, the neutrino oscillation processes generate classic NG fields that are strong enough in Supernovae to modify the neutrino flavor dynamics. The oscillation patterns may change in the periods of largest ν fluxes. Two examples are given: 1. the back reaction of a NG field improves the adiabaticity of the ν_e resonant conversion; 2. $\bar{\nu}_e \leftrightarrow \bar{\nu}_\mu$ oscillations may occur even if ν_e is the lightest of the neutrinos.

*Talk given at the International Symposium on Lepton and Baryon Number Violation, Trento, Italy, 1998

1 Basic Features

The Goldstone theorem asserts that in a theory with spontaneous breaking of a global symmetry group some massless scalar bosons should exist - the so-called Nambu-Goldstone bosons (NG) - one per global symmetry broken by the vacuum. Good candidates for spontaneously broken quantum numbers are the partial lepton numbers L_e, L_μ, L_τ and total lepton number L . Indeed, they are conserved by the Standard Model (SM) interactions, but several experiments going now from solar to atmospheric neutrinos and laboratory oscillation experiments [1] indicate that the lepton flavors are not conserved.

The NG bosons not only have zero mass but also no scalar potential terms such as ϕ^4 . Furthermore, they also interact with the fermion particles however, they do not mediate long range forces. The reason lies in the very global symmetry that, broken by the vacuum still operates at the Lagrangian level. A NG boson transforms as $\phi \rightarrow \phi + \alpha$, $\alpha = \text{constant}$, and therefore the symmetry requires that these fields have nothing but derivative couplings (this is related to the soft pion low energy theorems [2, 3]) *i.e.*, all of the NG boson interactions can be expressed with an effective Lagrangian of the form

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi + \mathcal{L}_{\text{int}}(\partial_\mu \phi). \quad (1)$$

The ϕ equation of motion is nothing but the conservation law of the total current associated with the $\phi \rightarrow \phi + \alpha$ symmetry,

$$\partial_\mu \partial^\mu \phi = -\partial_\mu J_\Lambda^\mu / V_\Lambda, \quad (2)$$

where V_Λ is essentially the scale of symmetry breaking and the current J_Λ^μ is determined in leading order by the quantum numbers of the particles. For instance, if Λ is the electron lepton number L_e ,

$$J_e^\mu = \bar{e} \gamma^\mu e + \bar{\nu}_e \gamma^\mu \nu_e + \dots, \quad (3)$$

where the dots stand for model dependent radiative corrections and scalar boson contributions.

This general result can be obtained using a suitable rephasing of the fields present in the theory [4, 5]. Let Λ_a and Λ_i be the quantum numbers of the fermion fields χ^a and scalars σ_i under an abelian global symmetry $U(1)_\Lambda$ which is spontaneously broken by vacuum expectation values $\langle \sigma_i \rangle$. A suitable change of variables namely,

$$\chi^a = e^{-i\Lambda_a \phi / V_\Lambda} \psi^a, \quad (4)$$

$$\sigma_i = e^{-i\Lambda_i \phi / V_\Lambda} (\langle \sigma_i \rangle + \rho_i), \quad (5)$$

makes the Lagrangian to be expressed in terms of *physical weak* eigenstates, fermions $\psi^a = e, \nu_e, \dots$, massive bosons ρ_i and gauge bosons (assumed to be singlets of the global symmetry) all of them invariant under $U(1)_\Lambda$. The only non-invariant field is the NG boson ϕ . It transforms as $\phi \rightarrow \phi + \alpha$ hence, the Lagrangian, symmetric under $U(1)_\Lambda$, can only depend on ϕ through the derivatives $\partial_\mu \phi$. Its leading order couplings are derived

from the kinetic Lagrangians of χ^a and σ_i by application of Eqs. (4-5). One obtains in particular, $V_\Lambda^2 = 2 \sum |\Lambda_i \langle \sigma_i \rangle|^2$.

The second member of the NG equation of motion, Eq. (2), is clearly not an ordinary scalar density. If one calculates the divergency of a vector or axial-vector current using the Dirac equation, one ends up with either scalar but flavor violating terms,

$$\partial_\mu \bar{\psi}_i \gamma^\mu \psi_j = i(m_i - m_j) \bar{\psi}_i \psi_j ,$$

or flavor conserving but pseudo-scalar,

$$\partial_\mu \bar{\psi}_i \gamma^\mu \gamma_5 \psi_j = i(m_i + m_j) \bar{\psi}_i \gamma_5 \psi_j .$$

This kind of source terms do not generate long range '1/r' fields, but rather spin dependent '1/r³' potentials [6].

In the intent of escaping the derivative couplings law, models were made with explicit symmetry breaking terms. These might be anomaly terms [7], as in the axion case [8], or soft symmetry breaking terms [9]. In this way scalar couplings with the NG field are produced but at the cost of giving a mass to the then, so-called pseudo Nambu-Goldstone boson. The mass ultimately means a finite range for the NG bosons. I will show next that under certain conditions a true Nambu-Goldstone boson may develop a macroscopic field [5, 10].

2 Long range Nambu-Goldstone fields

First notice that the source of a NG field is different from zero if the matter current J_Λ^μ is not conserved. The volume integral

$$\int d^3x \partial_\mu J_\Lambda^\mu = \frac{d\Lambda^{\text{created}}}{dt}$$

represents the rate of Λ -number *created* per unity of time (in all particles except ϕ) and $\partial_\mu J_\Lambda^\mu$ is its volume density. The equation of motion (2) gives

$$\phi(t, \vec{r}) = \frac{-1}{V_\Lambda} \int d^3x \frac{\dot{\rho}_\Lambda^{\text{cr}}(t - |\vec{r} - \vec{x}|, \vec{x})}{4\pi |\vec{r} - \vec{x}|} . \quad (6)$$

It is a long-range Coulomb type of field but suppressed by the product of the scale V_Λ times the interval of time per Λ -violating process. To have a significant macroscopic field one needs a large rate of reactions with a net Λ number variation. That calls for astrophysical sources. There is evidence that solar neutrinos change flavor as they come out of the Sun. If ν_e oscillates into ν_μ , for instance, then L_e and L_μ are not conserved in a large scale in stars and possibly in Supernovae as well. If any of them is associated with a spontaneously broken global symmetry then, the respective NG boson acquires classic field configurations.

For definiteness suppose that L_e is spontaneously broken and that a fraction of the ν_e s emitted from a star are resonantly converted into ν_μ through the Mikheyev-Smirnov-Wolfenstein (MSW) mechanism [11, 12] (also ν_μ into ν_e in a Supernova) in a certain shell of radius $\sim r_c$ inside the star. Assuming for simplicity stationary neutrino fluxes and spherical symmetry one obtains the solution for the NG boson ϕ_e :

$$\begin{aligned}\phi_e &= \text{const.} , \quad r < r_c , \\ \phi_e &= \frac{1}{V_\Lambda} \frac{1}{4\pi r} \left[\frac{dN}{dt}(\nu_e \rightarrow \nu_\mu) - \frac{dN}{dt}(\nu_\mu \rightarrow \nu_e) \right] , \quad r > r_c\end{aligned}\quad (7)$$

in terms of the total numbers of converted neutrinos, $N(\nu_\mu \rightarrow \nu_e)$ and $N(\nu_e \rightarrow \nu_\mu)$, per unity of time.

What are the possible consequences of such a NG field? Applying Eqs. (4) to the kinetic terms

$$\mathcal{L}_0 = \bar{e} i \not{\partial} e + \bar{\nu}_e i \not{\partial} \nu_e ,$$

one obtains the interaction

$$\mathcal{L}_{\text{int}} = \frac{1}{V_\Lambda} \partial_\mu \phi_e (\bar{e} \gamma^\mu e + \bar{\nu}_e \gamma^\mu \nu_e) . \quad (8)$$

The fermions only care about the gradient of the NG field. In the situation described above, denoting by $\vec{j}(\nu_e \rightarrow \nu_\mu)$ the flux of e -neutrinos converted to ν_μ and $\vec{j}(\nu_\mu \rightarrow \nu_e)$ the reciprocal,

$$\vec{A}_e = -\frac{\vec{\nabla} \phi_e}{V_\Lambda} = \frac{1}{V_\Lambda^2} [\vec{j}(\nu_e \rightarrow \nu_\mu) - \vec{j}(\nu_\mu \rightarrow \nu_e)] . \quad (9)$$

The quantity $\vec{A}_e$ is suppressed relatively to the field ϕ_e by the scale V_Λ and radius r and that is what makes it extremely small. However, as follows from Eq. (8), it still contributes to the potential energy of an electron-neutrino with velocity $\vec{v}_\nu$ as

$$V_{\nu_e} = \vec{A}_e \cdot \vec{v}_\nu . \quad (10)$$

These contributions associated with breaking of partial lepton numbers are flavor dependent by nature and therefore may have a role in neutrino oscillations.

One estimates the order of magnitude of theses NG potentials as the neutrino flux divided by V_Λ^2 . In the case of the Sun, evaluating the total flux at the Sun radius $R_\odot = 7 \times 10^{10}$ cm,

$$V_{\text{NG}} \sim \frac{j_\nu}{V_\Lambda^2} = \frac{2.8 \times 10^{-2} \text{ 1 keV}^2}{R_\odot V_\Lambda^2} .$$

It produces a too small variation of the neutrino phase, $\sim V_{\text{NG}} R_\odot$, thus it does not seem significant for solar neutrinos. In a Supernova however the fluxes are many orders of magnitude larger. One has

$$\frac{j_\nu}{V_\Lambda^2} = 1.48 \frac{G_\Lambda}{G_F} \frac{L_\nu}{10^{52} \text{ ergs/s}} \frac{10 \text{ MeV}}{\langle E_\nu \rangle} \left(\frac{r}{10^{10} \text{ cm}} \right)^{-2} \times 10^{-12} \text{ eV} , \quad (11)$$

where $G_F = 11.66 \text{ TeV}^{-2}$ is the Fermi constant, $G_e = 1/V_\Lambda^2$, L_ν is the neutrino energy luminosity, and $\langle E_\nu \rangle$ its average energy. $L_\nu / \langle E_\nu \rangle$ gives the neutrino emission rate. This has to be compared with the Standard Model electro-weak potential V_W [11]. In the regions of a Supernova star with densities typical of the Sun the mass density goes as $1/r^3$, the constant $\tilde{M} = \rho r^3$ lying between 10^{31}g and $15 \times 10^{31} \text{g}$ depending on the star [13]. Then,

$$V_W = \sqrt{2} G_F n_e = 0.76 Y_e \frac{\tilde{M}}{10^{31} \text{g}} \left(\frac{r}{10^{10} \text{cm}} \right)^{-3} \times 10^{-12} \text{ eV} , \quad (12)$$

$Y_e \approx 1/2$ being the electron abundance and n_e the number density. Clearly, in the period of time where the neutrino luminosities are high enough say, 10^{52} ergs/s , the scale of lepton symmetry breaking can be as high as the Fermi scale and yet the NG potentials compete with V_W at the places where the resonance occurs for values of Δm^2 interesting for solar neutrino solutions ($\Delta m^2/E \sim 10^{-12} \text{ eV}^2$). Furthermore, the NG potentials decay as $1/r^2$ and necessarily overcome the local interactions at large enough radius.

At this point we can already draw some conclusions. First, if the lepton numbers $L_e, L_\mu, \dots$ are spontaneously broken, the neutrino oscillations in a Supernova originate the appearance of classic Nambu-Goldstone fields. Second, if the scale of symmetry breaking lies below 1 TeV they in turn yield potential energies that are significant for the very neutrino oscillation processes. What are the possible consequences? The NG coupling are flavor dependent by nature hence, may alter the pattern of neutrino oscillations, with a fundamental difference: the NG potentials are proportional to the neutrino fluxes and that induces a time dependence of the effects in a Supernova context.

Next I show two examples. One [10] is the improvement of adiabaticity of otherwise $\nu_e \leftrightarrow \nu_X$ non-adiabatic transitions. The other [5] is the violation of the SM prediction that the anti-neutrinos do not resonantly oscillate if the neutrino masses obey the hierarchy of the charged lepton masses.

3 Improvement of adiabaticity: Majoron back reaction

Suppose that in the Sun the electron-neutrino oscillates into the muon-neutrino with the parameters of the non-adiabatic, small mixing angle solution [14, 15, 16]. Then, the $\nu_e \leftrightarrow \nu_\mu$ transitions are also non-adiabatic in a Supernova and only a fraction of each neutrino species is converted into the other. Furthermore, since the level crossing probability is an increasing function of the energy, the hotter ν_μ s have larger survival probabilities than the cooler ν_e s. Let us assume that the partial lepton number L_e is conserved at the Lagrangian level but the respective global symmetry, $U(1)_e$, is spontaneously broken. Then, a NG boson (Majoron), ϕ_e , exists with zero mass. Although in principle, the neutrino mass matrix violates all three lepton numbers, I will ignore for simplicity other

possibly existing Majorons. Notice that other NG fields are naturally less intense if the respective scales of symmetry breaking are slightly higher.

The equations governing two flavor oscillation [17] with mixing angle θ and Δm^2 difference in the vacuum, are given by, up to an universal phase absorbed in the wave functions,

$$i \frac{\partial}{\partial r} \begin{pmatrix} \nu_e \\ \nu_\mu \end{pmatrix} = \frac{1}{2E} \begin{pmatrix} 2E V_e & \frac{1}{2} \Delta m^2 \sin 2\theta \\ \frac{1}{2} \Delta m^2 \sin 2\theta & \Delta m^2 \cos 2\theta \end{pmatrix} \begin{pmatrix} \nu_e \\ \nu_\mu \end{pmatrix}. \quad (13)$$

V_e is the difference between the ν_e and ν_μ potentials, which is the following in the presence of a ϕ_e field (see Eqs. (10,12)):

$$V_e = V_{\nu_e} - V_{\nu_\mu} = V_W + \vec{A}_e \cdot \vec{v}_\nu. \quad (14)$$

Assuming spherical symmetry, $\vec{A}_e$ only has radial component and one obtains from Eq. (9)

$$V_e = \sqrt{2} G_F n_e + \frac{G_e}{4\pi r^2} [\dot{N}(\nu_e \rightarrow \nu_\mu) - \dot{N}(\nu_\mu \rightarrow \nu_e)]. \quad (15)$$

V_e would be the same if one considers alternatively a NG boson associated with the breaking of L_μ or $L_e - L_\mu$. The reason is, $\nu_e \leftrightarrow \nu_\mu$ oscillations only violate $L_e - L_\mu$ not the total number $L_e + L_\mu$.

In order to calculate the radial component $A_e(r)$ one needs to know the fluxes of ν_μ and ν_e as functions of the radius and their energy spectra. Let $f_{\nu_e}(E)$ and $f_{\nu_\mu}(E)$ be the number of neutrinos emitted per unity of time and energy, $d\dot{N}_{\nu_e}/dE$ and $d\dot{N}_{\nu_\mu}/dE$ respectively. If the transition is non-adiabatic, the number of converted neutrinos with a given energy E is only a fraction $1 - P_c$ of the radiated particles, P_c is the level crossing probability, and so

$$d\dot{N}(\nu_e \rightarrow \nu_\mu) = (1 - P_c) f_{\nu_e}(E) dE, \quad (16)$$

$$d\dot{N}(\nu_\mu \rightarrow \nu_e) = (1 - P_c) f_{\nu_\mu}(E) dE. \quad (17)$$

The problem of obtaining these rates as functions of the radius is simplified by neglecting the oscillation length of the resonantly converted neutrinos: in that limit a neutrino with energy E is converted to the other flavor at the position where the resonance condition is verified:

$$E(r) = \frac{\Delta m^2 \cos 2\theta}{2 V_e(r)}. \quad (18)$$

In addition, the level crossing probability P_c can be calculated using the Landau-Zener approximation [14, 17],

$$P_c(E) = \exp \left\{ -\frac{\pi}{4} \left| \frac{\Delta m^2 \sin^2 2\theta}{dE/dr \cos 2\theta} \right| \right\}_{\text{res.}}. \quad (19)$$




Figure 1: Level crossing probability as a function of the ν energy. Dashed curve: SM with $\tilde{M} = 4 \times 10^{31}$ g and $Y_e = 1/2$. Dotted and bold curves: a Nambu-Goldstone field exists with $G_e = G_F$ and $G_e = 4 G_F$ respectively. The luminosities are 10^{52} ergs/s for ν_e and 7×10^{51} ergs/s for ν_μ .

These approximations change the numbers but not, I believe, the lesson taken by comparing the results with and without a Majoron field. Eqs. (12,15 - 19) establish the differential equations for $d\dot{N}(\nu_e \rightarrow \nu_\mu)$ and $d\dot{N}(\nu_\mu \rightarrow \nu_e)$. They are non-linear because the derivative of $E(r)$ is not independent from the Majoron potential $A_e(r)$.

Let us examine the $\nu_e \leftrightarrow \nu_\mu$ oscillations with the mixing parameters of the non-adiabatic solar neutrino solution (for a recent update see [18]), choosing in particular the values $\Delta m^2 = 7 \times 10^{-6} \text{ eV}^2$, $\Delta m^2 \sin^2 2\theta = 4 \times 10^{-8} \text{ eV}^2$. In a Supernova the resonance is non-adiabatic as well and, as Eq. (19) indicates, the survival probability $1 - P_c$ increases with the ν energy. A detailed study in the framework of the SM was done in [19]. In Fig. 1, P_c in the Landau-Zener approximation is plotted against the energy. The dashed curve holds for the SM potential with a constant $\tilde{M} = 4 \times 10^{31}$ g. It is manifest the aggravation of the non-adiabaticity with the energy.

To study the Majoron case one has to specify the energy spectra and luminosities. I used Fermi-Dirac distributions with the following values of temperature and chemical potential [20]: for ν_e , $T = 2.4$ MeV and $\mu = 3.2 T$; for ν_μ , $T = 5.1$ MeV, $\mu = 4.1 T$. This gives average energies of 10 and 23 MeV respectively. The luminosity intensities are in turn 10^{52} ergs/s for ν_e and 7×10^{51} ergs/s for ν_μ which amount to particle emission rates of 10^{51} and 3×10^{50} ergs/s/MeV respectively. Because the e -neutrinos are more numerous, the $\nu_e \leftrightarrow \nu_\mu$ oscillations produce a net destruction of L_e -number and a positive potential A_e . The dynamics is the following: the less energetic ν_e oscillate to ν_μ at smaller radius




Figure 2: Total potential $V_e = V_{\nu_e} - V_{\nu_\mu}$ as a function of the radius. The dashed curve is the SM potential, the dotted and bold curves are the Majoron cases with the same parameters as in Fig. 1.

than $\nu_\mu \rightarrow \nu_e$; the $\nu_e \rightarrow \nu_\mu$ conversion produces a positive $A_e(r)$ which attenuates the fall of the total potential $V_e = V_W + A_e$ with r ; Consequently, the adiabaticity improves at larger radius and the most energetic neutrinos change flavor with higher probabilities.

Fig. 2 [10] shows the total potential V_e as a function r . The dashed curve stands for the SM potential, V_W , with $\tilde{M} = 4 \times 10^{31}$ g and $Y_e = 1/2$. The dotted and bold curves differ in the scale of L_e symmetry breaking, $G_e = G_F$ and $G_e = 4 G_F$, respectively. The potential falls more slowly as the field ϕ_e grows which reflects on the level crossing probability plotted in Fig. 1 [10] for the same cases (dotted and bold curves). The effect is clear: the stronger the Majoron field, the more efficient is the flavor conversion. It is worth to mention that if the μ -neutrinos were more numerous than the e -neutrinos the effect would be the opposite because the Majoron potential would be negative ($\dot{L}_e > 0$). That shows up in the rise of P_c at the ν_μ energy band around 20 MeV.

Fig. 3 [10] shows the implications for the outgoing ν_e energy spectrum. The dotted curves are the assumed luminosity distributions for the emitted ν_{eS} and $\nu_{\mu S}$. The dashed curve represents the luminosity of the e -neutrinos that come out of the star after standard MSW oscillations and the bold curve is the same but with a Majoron field ($G_e = 4 G_F$). The improvement of adiabaticity makes more $\nu_{\mu S}$ convert into ν_e and less ν_e to survive, and because the μ -neutrinos are more energetic, the outgoing ν_e spectrum is harder than if there was no Majoron field. The average energy of the outgoing e -neutrinos is 13 MeV for $G_e = 0$ but reaches 17 MeV if $G_e = G_F$ and 21 MeV if $G_e = 4 G_F$. These effects on the spectra can in principle be traced in those detectors such as Super-Kamiokande and SNO, capable of detecting Supernova electron-neutrinos [21].




Figure 3: The dotted curves are the assumed luminosity distributions for ν_e and ν_μ as emitted from the neutrinospheres. The dashed curve represents the ν_e luminosity after electroweak neutrino oscillations whereas in the bold curve a ϕ_e field exists with $G_e = 4 G_F$.

To summarize, if the explanation of the solar neutrino deficit is the MSW non-adiabatic oscillation $\nu_e \rightarrow \nu_\mu$ ($\nu_e \rightarrow \nu_\tau$ or any other superposition of ν_μ and ν_τ) then, the Standard Model predicts that in a Supernova the $\nu_e \leftrightarrow \nu_\mu$ transitions are also non-adiabatic. It means that, to a large extent, the e -neutrinos preserve their lower energy spectrum, unless ν_e also mixes to another flavor with a too high or too low Δm^2 to show up in solar neutrinos. If however, L_e is a spontaneously broken quantum number, the associated Nambu-Goldstone boson, ϕ_e , will acquire a classic field configuration which may be strong enough to produce a back reaction that improves the adiabaticity of the $\nu_e \leftrightarrow \nu_\mu$ transitions. The outcome is a ν_e energy spectrum harder than expected. In 1987, the existing detectors were only able to detect electron anti-neutrinos but the now operating Super-Kamiokande and SNO experiments will be capable of detecting Supernova ν_e events. The analysis of the energy distribution can in principle reveal or put limits on that kind of effect.

The scenarios of neutrino mixing change considerably if one considers the evidences from atmospheric and terrestrial neutrino experiments (for a review see [1]). The atmospheric neutrino anomaly and the zenith angle dependence observed by Super-Kamiokande can be explained by $\nu_\mu \rightarrow \nu_\tau$ oscillations[22], $\nu_\mu \rightarrow \nu_e$ excluded CHOOZ [23], but that is still consistent with solar neutrinos oscillations into a linear combination of ν_μ and ν_τ . That is no longer true if one takes in consideration the LSND evidences of $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ [24]

and $\nu_\mu \rightarrow \nu_e$ [25]. The much larger Δm^2 scale involved in LSND ($\Delta m_{e\mu}^2 > 0.2 \text{ eV}^2$), calls for a fourth flavor - a sterile neutrino - into which solar ν_e s should oscillate. That changes some features concerning the effects of a NG field ϕ_e but one thing is maintained: the signature of NG fields is a *surprise i.e.*, an oscillation pattern not consistent with SM weak interactions [10].

An important point is that the Nambu-Goldstone fields are proportional to the reaction rates that violate the relevant quantum numbers. In the case of neutrino oscillations this manifests as a dependence on the neutrino luminosity magnitudes. The effects of the Majoron fields if any, will be observed in a shorter or longer interval of time depending on the actual scale of lepton number symmetry breaking. Thus, the observation of such correlation with the flux intensities, would thus provide a measurement of the scale of spontaneous symmetry breaking.

4 Resonant conversion of anti-neutrinos

In the framework of the SM weak interactions the anti-neutrinos cannot undergo a resonant oscillation if the neutrino masses follow the same hierarchy as the charged leptons. That is because the SM potential V_W is positive in an electron rich star medium and the potential reverses sign for antineutrinos. The NG boson contributions are not only non-universal in flavour but also their signs depend on the flavour content of the neutrino fluxes and dominant flavor transitions. This allows in principle other than the standard MSW type of oscillations.

If one of the partial lepton numbers is spontaneously broken one expects the same happens for all three flavours. Let σ_i be the set of scalar fields, singlets of $SU(2) \times U(1)$, with quantum numbers $\Lambda_i = (L_e, L_\mu, L_\tau)_i$ under the symmetry group $U(1)_e \times U(1)_\mu \times U(1)_\tau$. If no residual symmetry is left after spontaneous symmetry breaking there are three NG bosons. By going to the physical basis as in Eqs. (4-5), now summing over the quantum numbers $\Lambda = L_e, L_\mu, L_\tau$,

$$\chi^a = e^{-i \sum \xi_\Lambda \Lambda_a} \psi^a, \quad (20)$$

$$\sigma_i = e^{-i \sum \xi_\Lambda \Lambda_i} (\langle \sigma_i \rangle + \rho_i), \quad (21)$$

one obtains the Lagrangian of the NG fields

$$\mathcal{L} = \frac{1}{2} V_{\Lambda M}^2 \partial_\mu \xi_\Lambda \partial^\mu \xi_M + \mathcal{L}_{\text{int}}(\partial_\mu \xi_\Lambda), \quad (22)$$

the sum over Λ, M is implicit. The matrix

$$V_{\Lambda M}^2 = 2 \sum_i \Lambda_i M_i |\langle \sigma_i \rangle|^2 \quad (23)$$

is in general non-diagonal, provided that some of the fields σ_i have simultaneously more than one non-trivial quantum number. Such mixing implies that the NG bosons mediate

an action at a distance between two distinct currents as show the equations of motion,

$$\partial_\mu \partial^\mu \xi_\Lambda = - \sum_M G_{\Lambda M} \partial_\mu J_M^\mu, \quad (24)$$

where $G_{\Lambda M}$ is the inverse matrix of $V_{\Lambda M}^2$.

In a stationary regime, if the neutrinos oscillate causing current variations $\delta \vec{J}_M$, the expression for the gradients of the NG bosons, $\vec{A}_\Lambda = -\vec{\nabla} \xi_\Lambda$, that generalizes Eq. (9) is [5]:

$$\vec{A}_\Lambda = - \sum_M G_{\Lambda M} \delta \vec{J}_M. \quad (25)$$

Since the interaction Lagrangian of the leptons has the same form as Eq. (8) *i.e.*, $\mathcal{L}_{\text{int}} = J_\Lambda^\mu \partial_\mu \xi_\Lambda$, the potential energy of a neutrino ν_ℓ subject to stationary NG fields is given by

$$V_{\nu_\ell} = \vec{A}_\ell \cdot \vec{v}_\nu, \quad (26)$$

as in Eq. (10). Plugging in Eq. (25), one gets for example,

$$V_{\nu_\mu} = - \left(G_{\mu e} \delta \vec{J}_e + G_{\mu\mu} \delta \vec{J}_\mu + G_{\mu\tau} \delta \vec{J}_\tau \right) \cdot \vec{v}_\nu.$$

This shows that the violation of a certain lepton number in one place originates fields that acts on other flavors in other places.

In the old preferred scenario (before Super-Kamikande results on atmospheric neutrinos [22]), the neutrino mixing angles are small, $\nu_e \approx \nu_1$, $\nu_\mu \approx \nu_2$, $\nu_\tau \approx \nu_3$, and the masses obey the hierarchy $m_1 < m_2 < m_3$. Then [26, 27], as the neutrinos come out of a Supernova the first resonant transitions are between ν_e and ν_τ . It is outside their resonance shell, at lower medium densities, that MSW transitions $\nu_e \leftrightarrow \nu_\mu$, with smaller Δm^2 , occur.

In this meeting Atsuto Suzuki [22] reported Super-Kamiokande results on the atmospheric neutrino anomaly and zenith angle dependence that can be explained by $\nu_\mu \rightarrow \nu_\tau$ oscillations. The best values, confirmed at Neutrino 98, are $\Delta m^2 = 5 \times 10^{-3} \text{ eV}^2$ and $\sin^2 2\theta = 1$. The alternative $\nu_\mu \rightarrow \nu_e$ oscillation is excluded by CHOOZ [23]. This is consistent with a scenario where the heaviest of the neutrinos is essentially a superposition, ν_H (possibly maximal), of ν_μ and ν_τ ; ν_e is the lightest neutrino and oscillates in the Sun into the medium weight orthogonal superposition, ν_L , of ν_μ and ν_τ . In what follows what is said about ν_τ , ν_μ and quantum numbers L_τ , L_μ can be maintained if replaced by ν_H , ν_L and respective combinations of L_τ and L_μ .

In a Supernova the ν_e flux is larger than the ν_τ flux and consequently more ν_e transform into ν_τ than the reverse which makes a net decrease of $L_e - L_\tau$. The current densities $\vec{J}_\tau$ and $\vec{J}_e$ suffer the variations

$$\delta \vec{J}_\tau = -\delta \vec{J}_e = \vec{j}(\nu_e \rightarrow \nu_\tau) - \vec{j}(\nu_\tau \rightarrow \nu_e), \quad (27)$$

that generate the NG fields ξ_e , ξ_μ , ξ_τ . Applying Eq. (25) one gets the vector potentials outside the sphere where $\nu_e \leftrightarrow \nu_\tau$ takes place. The magnitude of the constants $G_{\Lambda M}$

is directly related to the symmetry breaking scales, but the flavor structure is arbitrary and is not constrained by the neutrino mixing. The matrix $G_{\Lambda M}$ has only to be symmetric and positive definite as demands the scalar bosons kinetic Lagrangian. Consider the following hierarchy: $|G_{ee}|, |G_{e\mu}|, |G_{e\tau}| \ll |G_{\mu\tau}|, G_{\tau\tau}$. Then, $\vec{A}_e$ is much smaller in magnitude than $\vec{A}_\mu$ and $\vec{A}_\tau$ and, assuming spherical symmetry, the potential energies are well approximated by

$$V_{\nu_e} = V_W + A_e, \quad (28)$$

$$V_{\nu_\mu} = A_\mu \approx -G_{\mu\tau} \delta J_\tau, \quad (29)$$

$$V_{\nu_\tau} = A_\tau \approx -G_{\tau\tau} \delta J_\tau, \quad (30)$$

omitted the neutral current terms. The anti-neutrino potentials are the symmetric of these.

The increment of τ lepton number per unity of time, $\delta F_\tau = \dot{N}(\nu_e \rightarrow \nu_\tau) - \dot{N}(\nu_\tau \rightarrow \nu_e)$, is positive and so is the flux transfer $\delta J_\tau = \delta F_\tau / 4\pi r^2$. Defining $\delta F_\tau = \delta F_{51} \times 10^{51} \text{ erg/s/MeV}$ one estimates $\delta F_{51} = 2-5$ in the first 1/2 seconds of neutrino emission until the Supernova explosion and $\delta F_{51} = 0.2-0.5$ at 1 second time. See [5] for more details. The NG potential for ν_τ is negative but that is not necessarily the case of ν_μ because the element $G_{\mu\tau}$ can be negative! Using Eq. (11),

$$A_\mu = -1.48 \frac{G_{\mu\tau}^F}{G_F} \delta F_{51} \left(\frac{r}{10^{10} \text{ cm}} \right)^{-2} \times 10^{-12} \text{ eV}. \quad (31)$$

The Schrödinger equation for the anti-neutrinos is the same as for neutrinos, Eq. (13), but with a symmetric potential:

$$i \frac{\partial}{\partial r} \begin{pmatrix} \bar{\nu}_e \\ \bar{\nu}_\mu \end{pmatrix} = \frac{1}{2E} \begin{pmatrix} -2E(V_W - A_\mu) & \frac{1}{2}\Delta m^2 \sin 2\theta \\ \frac{1}{2}\Delta m^2 \sin 2\theta & \Delta m^2 \cos 2\theta \end{pmatrix} \begin{pmatrix} \bar{\nu}_e \\ \bar{\nu}_\mu \end{pmatrix}, \quad (32)$$

where $|A_e| \ll |A_\mu|$ is implicit. Within the SM, the effective mass $-2E V_W$ would be negative, and the resonance condition $-2E V_W = \Delta m^2 \cos 2\theta$ could not be met. In presence of NG fields, the effective mass $M_e^2 = -2E(V_W - A_\mu)$ is negative at the high density regions, but since the NG potentials only decay with the second power of the radius their contributions inevitably overcome at large enough distances. That is a crucial point. If on top of that $G_{\mu\tau}$ is negative, the effective mass M_e^2 suddenly turns positive and may reach the value $M_\mu^2 = \Delta m^2 \cos 2\theta$. In that case there is a resonance layer where $M_e^2 = M_\mu^2$ is met. In Fig. 4 it corresponds to the interception of the "e" and "μ" curves, $M_e^2(r)$ and M_μ^2 , respectively. The positive part of the 'potential' M_e^2 is plotted in Fig. 4 with values $G_{\mu\tau} = -G_F$, $\delta F_{51} = 1$ or 3, $Y_e = 1/2$, $\tilde{M}_{31} = 4$, and a neutrino energy of 20 MeV, the same for all curves.

Take $M_\mu^2 = 6 \times 10^{-7} \text{ eV}^2$ for example. The M_μ^2 dashed line crosses the relative potential curve at two points: the first, r_1 , lies in the remarkably sharp transition from the $-V_W < 0$ dominated region to the NG field dominated one and very likely a non-adiabatic level




Figure 4: $2E(V_{\bar{\nu}_e} - V_{\bar{\nu}_\mu})$ is plotted in the solid and dotted curves with $\delta F_{51} = 1$ and 3 respectively. The dashed line stands for a particular value of $\Delta m^2 \cos 2\theta$ and the squares curve is the adiabatic boundary for $M_\mu^2 \tan^2 2\theta$ in resonance 2.

crossing occurs there *i.e.*, $\bar{\nu}_e$ continues $\bar{\nu}_e$ and $\bar{\nu}_\mu$ continues $\bar{\nu}_\mu$. In the second point, r_2 , the potential is smooth enough ($M_e^2 \propto r^{-2}$) for adiabatic transitions. In general terms, the conversion probability is enhanced [12, 17] if $M_\mu^2 \tan^2 2\theta$ is larger than the rate $f(r) = |E dV/V dr|$ at resonance. In Fig. 4, $f(r)$ is plotted with squares for $V \propto r^{-2}$. One can see that small mixing angle adiabatic conversions are possible for a wide range of neutrino masses (for more details see [5]). Through this mechanism one could have resonant $\bar{\nu}_e \leftrightarrow \bar{\nu}_\mu$ oscillations in a Supernova whereas $\nu_e \rightarrow \nu_\mu$ take place in the Sun. In particular, with the parameters of the non-adiabatic solution [14, 15, 16] namely, $M_\mu^2 \approx (3 - 12) \times 10^{-6} \text{ eV}^2$ and $M_\mu^2 \tan^2 2\theta \approx 4 \times 10^{-8} \text{ eV}^2$, provided that the neutrino fluxes are high enough (compare the dotted and solid curves).

Since the predicted energy spectrum of $\bar{\nu}_\mu$ and $\bar{\nu}_\tau$ is harder than the $\bar{\nu}_e$ spectrum that kind of oscillation could be manifest in the SN1987A events [28]. Some analyses made [29, 30], but not all [31], disfavour the occurrence of $\bar{\nu}_e \leftrightarrow \bar{\nu}_\mu$ oscillations. This deserves some discussion [5], here I simply stress that the effects caused by the NG fields strongly depend on magnitude of the neutrino fluxes and on the scale of the global symmetry breaking as well. Since the fluxes vary with time in a Supernova, one could have anti-neutrino oscillations in the first instants of emission that fade away when the fluxes drop below a certain threshold. The observation in the future of such correlation between oscillation and flux magnitudes would provide a measurement of the symmetry breaking scale.

5 Conclusions

We have shown that despite the fact that true Nambu-Goldstone (NG) bosons only have derivative couplings, it is still possible to have coherent NG fields. This happens whenever the quantum numbers they are associated with are not conserved in reactions taking place in a large scale. If the lepton numbers are spontaneously broken charges then, neutrinos oscillations in stars produce classic NG fields. In addition, if the scale of symmetry breaking is below 1 TeV, these fields are strong enough in Supernovae to change the neutrino flavor dynamics.

The observable signatures are:

- Time variation of neutrino oscillation patterns in correlation with the flux magnitudes. This is a consequence of the NG fields being proportional to the time rate of reactions and therefore to the neutrino fluxes.
- Incongruence of Supernova neutrino oscillation patterns with the evidence obtained from solar, atmospheric and terrestrial neutrinos. The NG fields are not expected to be significant in these cases even if they are in Supernovae because the fluxes are much smaller. This is ultimately a result of the derivative coupling nature of the NG bosons.

Finally, if we will be lucky enough to see such kind of phenomena in the future, then, Supernova neutrinos will provide a measurement of the scale of lepton number symmetry breaking.

Acknowledgements

This work was supported in part by the project ESO/P/PRO/1127/96.

References

- [1] D. O. Caldwell, in Proceedings of "COSMO 97: First International Workshop on Particle Physics and the Early Universe", Ambleside, England, 1997 (edited by L. Roszkowski, World Scientific, 1998), hep-ph/9804367; Y. Totsuka, in Proceedings of the "XVIII International Symposium on Lepton-Photon Interactions", Hamburg, 1997 (edited by A. De Roeck and A. Wagner, World Scientific, 1998).
- [2] T.-P. Cheng and L.-F. Li, "Gauge Theory of Elementary Particle Physics" (Oxford University Press, New York, 1989).
- [3] S. Weinberg, "The Quantum Theory of Fields: II" (Cambridge University Press, Cambridge, England, 1996).
- [4] G. B. Gelmini, S. Nussinov and T. Yanagida, Nucl. Phys. B 219 (1983) 31.
- [5] L. Bento, Phys. Rev. D 57 (1998) 583.
- [6] Y. Chikashige, R. N. Mohapatra and R. D. Peccei, Phys. Lett. B 98 (1981) 265.
- [7] F. Wilczek, Phys. Rev. Lett. 49 (1982) 1549; D. Chang, R. N. Mohapatra and S. Nussinov, Phys. Rev. Lett. 55 (1985) 2835.
- [8] R. Peccei and H. Quinn, Phys. Rev. Lett. 38 (1977) 1440; S. Weinberg, Phys. Rev. Lett. 40 (1978) 223; F. Wilczek, Phys. Rev. Lett. 40 (1978) 279.
- [9] C. T. Hill and G. G. Ross, Nucl. Phys. B 311 (1988) 253; J. A. Frieman, C. Hill and R. Watkins, Phys. Rev. D 46 (1992) 1226; R. N. Mohapatra and S. Nussinov, Phys. Lett. B 395 (1997) 63.
- [10] L. Bento, hep-ph/9806411, Phys. Rev. D 59, 015013 (1999).
- [11] L. Wolfenstein, Phys. Rev. D 17 (1978) 2369.
- [12] S. P. Mikheyev and Y. Yu. Smirnov, Yad. Fiz. 42 (1985) 1441 [Sov. J. Nucl. Phys. 42 (1985) 913] and Nuovo Cimento 9C (1986) 17.
- [13] J. R. Wilson *et. al.*, Ann. N. Y. Acad. Sci. 470 (1986) 267.
- [14] W. C. Haxton, Phys. Rev. Lett. 57 (1986) 1271.
- [15] S. J. Parke, Phys. Rev. Lett. 57 (1986) 1275.
- [16] S. P. Rosen and J. M. Gelb, Phys. Rev. D 34 (1986) 969.
- [17] T. K. Kuo and J. Pantaleone, Rev. Mod. Phys. 61 (1989) 937.
- [18] N. Hata and P. Langacker, Phys. Rev. D 56 (1997) 6107.

- [19] H. Minakata and H. Nunokawa, Phys. Rev. D 38 (1988) 3605.
- [20] A. Burrows, Annu. Rev. Nucl. Sci. 40 (1990) 181.
- [21] A. Burrows, D. Klein, and R. Gandhi, Phys. Rev. D 45 (1992) 3361.
- [22] A. Suzuki (Super-Kamiokande collaboration), talk given at the 'International Symposium on Lepton and Baryon Number Violation', Trento, 1998.
- [23] M. Appollonio *et al.*, Phys. Lett. B. 420 (1998) 397.
- [24] C. Athanassopoulos *et al.*, Phys. Rev C 54 (1996) 2685; Phys. Rev. Lett. 77(1996) 3082.
- [25] C. Athanassopoulos *et al.*, Phys. Rev C 58 (1998) 2489; Phys. Rev. Lett. 81 (1998) 1774.
- [26] H. A. Bethe, Phys. Rev. Lett. 56 (1986) 1305.
- [27] T. P. Walker and D. N. Schramm, Phys. Lett. B 195 (1987) 331.
- [28] K. Hirata *et al.*, Phys. Rev. Lett. 58 (1987) 1490; R. M. Bionta, *ibid*, 1494.
- [29] B. Jegerlehner, F. Neubig and G. Raffelt, Phys. Rev. D 54 (1996) 1194.
- [30] A. Yu. Smirnov, D. N. Spergel and J. N. Bahcall, Phys. Rev. D 49 (1994) 1389.
- [31] P. J. Kernan and L. M. Krauss, Nucl. Phys. B 437 (1995) 243.